

PARC NATIONAL DE L'UPEMBA
I. MISSION G. F. DE WITTE
en collaboration avec
W. ADAM, A. JANSSENS, L. VAN MEEL
et R. VERHEYEN (1946-1949).
Fascicule 8 (4)

NATIONAAL UPEMBA PARK
I. ZENDING G. F. DE WITTE
met medewerking van
W. ADAM, A. JANSSENS, L. VAN MEEL
en R. VERHEYEN (1946-1949).
Aflevering 8 (4)

SCOLYTOIDEA **(COLEOPTERA)**

PAR

K. E. SCHEDL (Lienz) (*)

Le Président de l'Institut des Parcs Nationaux du Congo Belge m'a soumis une intéressante collection recueillie par la Mission d'exploration du Parc National de l'Upemba (1947-1949), sous la direction de M. G. F. DE WITTE.

La plupart des exemplaires ont été récoltés à la lumière; cependant, certains furent obtenus par tamisage de détritus ou de terre; d'autres dans des nids de rat-taupe, quelques-uns sous des feuilles mortes.

Ceci est très intéressant et prouve que quelques *Scolytidæ* congolais passent une partie de leur vie hors de leur plante-hôte, dans les débris au sol ou en terre, tout comme certaines espèces paléarctiques.

Cette diapause est certainement liée aux saisons sèche et humide.

La présence de quelques espèces dans des nids de rat-taupe est sans importance particulière, celles-ci étant certainement des trogloxènes.

L'habitude de passer une partie de leur vie au sol est vraisemblablement assez fréquente. Les exemplaires trouvés dans ces conditions appartiennent à neuf espèces et six genres, ressortissant à trois sous-familles. Ce sont :

Hypothenemus tuberculosus HAGEDORN.
Hypothenemus bauhaniae SCHEDL.

(*) 130. Contribution à la systématique et morphologie des *Scolytoidea*.

Stephanoderes polyphagus EGGERS.
Coccotrypes subovalis EGGERS.
Premnobius cavipennis EICHHOFF.
Premnobius adjunctus EGGERS.
Eccoptopterus sexspinosis MOTSCHULSKY.
Xyleborus principalis EICHHOFF.

Seules des études plus approfondies pourront établir jusqu'à quel point la connaissance de ce détail pourra servir dans la lutte contre les xylophages congolais.

CATALOGUE DES RÉCOLTES.

A. — IPIDÆ.

CTONOXYLON HAGEDORN.

Ctonoxylon HAGEDORN, 1910, Deutsche Ent. Zeitschr., p. 4.

1. — **Ctonoxylon setifer** EGGERS.

Ctonoxylon setifer EGGERS, 1920, Ent. Bl., XVI, p. 39.

Riv. Buye-Bala (affl. g. Muye, sous-affl. dr. Lufira), alt. 1.750 m, 1-7.IV.1948.

POLYGRAPHUS ERICHSON.

Polygraphus ERICHSON, 1836, Arch. Naturg., II, p. 57.

2. — **Polygraphus bicolor** EGGERS.

Polygraphus bicolor EGGERS, 1935, Rev. Zool. Bot. Afr., vol. XXVII, pp. 302-303.

Riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.XI.1947.

HYPOTHENEMUS WESTWOOD.

Hypothenemus WESTWOOD, 1836, Trans. Ent. Soc. London, I, p. 34.

3. — **Hypothenemus tuberculatus** HAGEDORN.

Hypothenemus tuberculatus HAGEDORN, 1912, Rev. Zool. Bot. Afr., vol. I, pp. 339-340.

Mabwe (rive Est lac Upemba), alt. 585 m, 3.I.1949; 21-24.II.1949 (recueillis à l'appareil de BERLESE, dans feuilles mortes).

4. — **Hypothenemus bauhaniæ** SCHEDL.

Hypothenemus bauhaniæ SCHEDL, 1950, Bull. Inst. roy. Sci. nat. Belg., XXVI, pp. 19-20.

Kabwe (sur rive dr. Muye, affl. dr. Lufira), alt. 1.320 m, 14.V.1948 (recueilli par tamisage); Mabwe (rive Est lac Upemba), alt. 585 m, 21-24.II.1949 (dans feuilles mortes).

STEPHANODERES EICHHOFF.

Stephanoderes EICHHOFF, 1871, Berl. Ent. Zeitschr., XV, p. 132.

5. — **Stephanoderes polyphagus** EGGERS.

Stephanoderes polyphagus EGGERS, 1924, Ent. Bl., vol. XX, p. 104.

Mabwe (rive Est lac Upemba), alt. 585 m, 21-24.II.1949 (dans feuilles mortes).

COCCOTRYPES EICHHOFF.

Coccotrypes EICHHOFF, 1878, Stett. Ent. Zeit., vol. XXXIX, p. 391.

6. — **Coccotrypes subovalis** EGGER.

Coccotrypes subovalis EGGER, 1932, Rev. Zool. Bot. Afr., vol. XXII, p. 291.

Kabwe (rive dr. Muye, affl. dr. Lufira), alt. 1.320 m, 12.V.1948.

MIMIPS EGGER.

Mimips EGGER, 1932, Rev. Zool. Bot. Afr., vol. 22, p. 33.

7. — **Mimips pilosus** EGGER.

Ips pilosus EGGER, 1924, Ent. Bl., vol. XX, p. 106.

Kaswabilenga (sur rive dr. Lufira), alt. 700 m, 16.X.1947.

PREMNObIUS EICHHOFF.

Premnobius EICHHOFF, 1879, Ratio Tomicin., p. 404.

8. — **Premnobius cavipennis** EICHHOFF.

Premnobius cavipennis EICHHOFF, 1879, Ratio Tomicin., p. 404.

Mukana (marais près de Lusinga), alt. 1.810 m, 15.III.1947; Lusinga, alt. 1.760 m, 21-26.III.1947 (un spécimen plus grand que ceux de la plaine);

riv. Kenia (affl. dr. Lusinga, sous-affl. dr. Lufwa), alt. 1.585 m, 29.II.1947 (dans nid de rat-taupe, spécimens plus grands que ceux de la plaine); gorges de la Pelenge, alt. 1.250 m, 31.V.1947; Mabwe (rive Est lac Upemba), alt. 585 m, 1-15.VII.1947; riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947; Kaswabilenga (rive dr. Lufira), alt. 700 m, 1-26.X.1947; riv. Lukawa, affl. dr. Lufira), alt. 700 m, 30.IX.1947; Kabwe (sur rive dr. Muye, affl. Lubanga), alt. 1.320 m, 12-14.V.1948 (dans terre tamisée).

9. — **Premnobius adjunctus** EGGERS.

Xyleborus adjunctus EGGERS, 1924, Ent. Bl., vol. XX, p. 108.

Lusinga, alt. 1.760 m, 31.II.1947; région confl. Mubale-Munte, alt. 1.480 m (dans nid de rat-taupe), 14.V.1947.

10. — **Premnobius minor** EGGERS.

Premnobius minor EGGERS, 1927, Rev. Zool. Bot. Afr., vol. XV, p. 183.

Mabwe (rive Est lac Upemba), alt. 585 m, 11.I.1949.

ECCOPTOPTERUS MOTSCHULSKY.

Eccoptopterus MOTSCHULSKY, 1863, Bull. Mosc., I, p. 515.

11. — **Eccoptopterus sexspinosis** MOTSCHULSKY.

Eccoptopterus sexspinosis MOTSCHULSKY, 1863, Bull. Mosc., I, p. 515.

Kabwe (sur rive dr. Muye, affl. Lufira), alt. 1.320 m, 12.V.1948 (recueilli par tamisage).

XYLEBORUS EICHHOFF.

Xyleborus EICHHOFF, 1864, Berl. Ent. Zeitschr., vol. VIII, p. 37.

12. — **Xyleborus ricini** EGGERS.

Xyleborus ricini EGGERS, 1932, Rev. Zool. Bot. Afr., vol. XXII, fasc. 3, p. 298.

Riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947.

13. — **Xyleborus forficulus** EGGER.

Xyleborus forficulus EGGER, 1922, Ent. Blätter, vol. XVIII, pp. 171-172.

Gorges de la Pelenge, alt. 1.250 m, 31.V.1947.

14. — **Xyleborus principalis** EICHHOFF.

Xyleborus principalis EICHHOFF, 1879, Rat. Tomin., p. 357.

Riv. Kenia (affl. dr. Lusinga, sous-affl. dr. Lufwa), alt. 1.585 m, 29.II.1948 (dans nid de rat-taupe).

15. — **Xyleborus badius** EICHHOFF.

Xyleborus badius EICHHOFF, 1868, Berl. Ent. Zeitschr., XII, p. 280.

Riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947; riv. Lukawe (affl. dr. Lufira), alt. 700 m, 30.IX au 22.X.1947; Kaswabilenga (rive dr. Lufira), alt. 700 m, 1.X au 4.XI.1947.

16. — **Xyleborus mascarensis** EICHHOFF.

Xyleborus affinis var. *B. mascarensis* EICHHOFF, 1879, Rat. Tom., p. 372.

Lusinga, alt. 1.760 m, 22.III.1947; 16.IV.1947; riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947; riv. Lukawe (affl. rive dr. Lufira), alt. 700 m, 30.IX au 22.X.1947; Kaswabilenga, alt. 700 m, 17.X.1947; 24.IX au 8.XI.1947; riv. Lupiala, alt. 850 m, 24.X.1947.

17. — **Xyleborus ferrugineus** FABRICIUS.

Xyleborus ferrugineus FABRICIUS, 1801, Syst. Eleuth., II, p. 388.

Lusinga, alt. 1.760 m, 12.IV.1947; riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947; riv. Lukawe (affl. dr. Lufira), alt. 700 m, 30.IX au 22.X.1947; riv. Lupiala (affl. dr. Lufira), alt. 700-850 m, 1-24.X.1947; riv. Mongolo (affl. g. Lufira), alt. 850 m, 28.X.1947; Kaswabilenga (rive dr. Lufira), alt. 700 m, 1.X au 8.XI.1947.

B. — PLATYPODIDÆ.

PLATYPUS HERBST.

Platypus HERBST, 1793, Natursyst., V, p. 128.

1. — **Platypus hintzi** SCHAUFFUS.

Platypus hintzi SCHAUFFUS, 1897, Berl. Ent. Zeitschr., XLII, p. 103.

Mabwe (rive Est lac Upemba), alt. 585 m, 1-15.VIII.1947; 6.II.1949 (recueillis à l'appareil de BERLESE); riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947; riv. Lukawe (affl. dr. Lufira), alt. 700 m, 30.IX au 22.X.1947; piste de Lusinga, alt. 1.200 m, 24.X.1947; riv. Lupiala, alt. 850 m, 24.X.1947; Kaswabilenga (rive dr. Lufira), alt. 700 m, 1.X au 8.XI.1947.

2. — **Platypus solutus** SCHEDL.

Platypus solutus SCHEDL, 1933, Rev. Zool. Bot. Afr., vol. XXIII, fasc. 3, pp. 193-194.

Riv. Lukawe (affl. dr. Lufira), alt. 700 m, 30.IX.1947; Kaswabilenga (rive dr. Lufira) alt. 700 m, 1-9.X.1947.

TRIOZASTUS SCHEDL.

Triozastus SCHEDL, 1939, Verh. 7 Int. Kongr. Ent. Berlin, I, p. 403.

3. — **Triozastus propatulus** SCHEDL.

Crossotarsus propatulus SCHEDL, 1935, Ent. Nachr. Bl., 9, pp. 153-154.

Kaswabilenga (rive dr. Lufira), alt. 700 m, 1.X au 8.XI.1947; riv. Lukawe (affl. dr. Lufira), alt. 700 m, 30.IX au 9.X.1947.

DOLIOPYGUS SCHEDL.

Doliopygus SCHEDL, 1939, Verh. 7 Int. Kongr. Ent. Berlin, I, p. 403.

4. — **Doliopygus serratus** STROHMEYER.

Crossotarsus serratus STROHMEYER, 1911, Ent. Blätt., vol. VII, pp. 226-227.

Mabwe (rive Est lac Upemba), alt. 585 m, 26.VIII.1947; riv. Lukawe (affl. dr. Lufira), alt. 700 m, 22.X.1947.

5. — **Doliopygus dubius** SAMPSON.

Crossotarsus crinitus var. *dubius* SAMPSON, 1924, Rev. Zool. Bot. Afr., vol. XII, p. 126.

Crossotarsus serratus var. STROHMEYER, 1911, Ent Blätt., vol. VII, p. 227.

Riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947.

6. — **Doliopygus opifex** SAMPSON.

Crossotarsus opifex SAMPSON, 1922, Ann. Mag. Nat. Hist., vol. IX, (9), pp. 139-140.

Lusinga, alt. 700 m, 26.III.1947; riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m. 25.IX.1947.

7. — **Doliopygus mimicus** SCHEDL.

Crossotarsus mimicus SCHEDL, 1933, Rev. Zool. Bot. Afr., vol. XXIII, fasc. 2, pp. 198-199.

Riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947; Lukawe (affl. dr. Lufira), alt. 700 m, 30.IX au 9.X.1947; Kaswabilenga (rive dr. Lufira), alt. 700 m, 1.X au 8.XI.1947.

8. — **Doliopygus præclarus** SCHEDL.

Crossotarssus præclarus SCHEDL, 1933, Rev. Zool. Bot. Afr., vol. XXIII, fasc. 2, pp. 199-200 (σ); SCHEDL, 1936, Rev. Zool. Bot. Afr., vol. XXIX, fasc. 1, p. 130 (φ).

Riv. Kande (affl. g. Lupiala, sous-affl. dr. Lufira), alt. 700 m, 25.IX.1947; Kaswabilenga (rive dr. Lufira), 700 m, 1-9.X.1947; riv. Lupiala (affl. dr. Lufira), alt. 700 m, 6-9.X.1947.

9. — **Doliopygus eximus** SCHEDL.

Doliopygus eximus SCHEDL, 1941, Rev. Zool. Bot. Afr., vol. XXXIV, p. 407.

Gorges de la Pelenge, alt. 1.150 m, 28.V.1947.

10. — **Doliopygus erichsoni** CHAPUIS.

Doliopygus erichsoni CHAPUIS, 1865, Monogr. d. Platyp., p. 95.

Mabwe (rive Est lac Upemba), alt. 585 m, 1-12.VIII.1947.

Doliopygus wittei n. sp.

Mâle. — Corps d'un brun rougeâtre; 2,77 fois aussi long que large. Cette espèce est unique dans le groupe des *Doliopygi abdominali* par la forme de l'échancrure médiane du deuxième sternite abdominal. Longueur : 4,9 mm.

Front plan; sillon médian extrêmement petit; surface mate, à ponctuation aréolée vers le vertex et sur les côtés, la ponctuation plus serrée aux angles latéro-antérieurs; bord de l'épistome brillant et lisse, peu ponctué; pubescence très courte et fine; séparation du vertex et du front anguleuse, mais pas très prononcée.

Pronotum à peine plus long que large (48 : 46), carré, les échancrures latérales assez profondes; surface lisse et brillante, très finement ponctuée, les points inégaux en grandeur et densité; quelques points pilifères le long du bord antérieur, quelques-uns plus gros vers la base.

Élytres distinctement plus larges que le pronotum (53 : 46) et à peu près deux fois aussi longs que larges, lisses et brillants, très régulièrement ponctués-striés, les points assez denses, les intervalles peu convexes, avec quelques points très fins et très épais, un peu plus nombreux à la base; stries plus enfoncées vers l'arrière; intervalles alternes aigus, comme d'habitude chez le groupe des *Doliopygi abdominalis*; abdomen avec les prolongements latéraux du second sternite très longs, l'échancrure médiane très grande,

de forme rectangulaire, triangulairement incisé au milieu, l'intérieur des prolongements très densément et fortement ponctué, hérissé de longs poils; avant de l'échancrure médiane pourvu d'une gouttière très large, formant un liteau de chaque côté; sternites 3 et 4 étroits, lisses et imponctués, sternite 5 concave, avec de gros points.

T y p e s : à l'Institut des Parcs Nationaux du Congo Belge et dans la collection SCHEDL.

L o c a l i t é s : Kaswabilenga, alt. 700 m, 6-7.XI et 17.X.1947.

Je dédie cette espèce nouvelle à M. G. F. DE WITTE, Conservateur à l'Institut royal des Sciences naturelles de Belgique, Chef de la Mission d'exploration du Parc National de l'Upemba.

PARC NATIONAL DE L'UPEMBA

I. MISSION G. F. DE WITTE

en collaboration avec

W. ADAM, A. JANSSENS, L. VAN MEEL

et R. VERHEYEN (1946-1949).

Fascicule 8 (5)

NATIONAAL UPEMBA PARK

I. ZENDING G. F. DE WITTE

met medewerking van

W. ADAM, A. JANSSENS, L. VAN MEEL

en R. VERHEYEN (1946-1949).

Aflevering 8 (5)

BIBIONIDÆ AND DORILAIDÆ (DIPTERA)

BY

D. ELMO HARDY (Honolulu, Hawaii)

Flies of these families were not well represented in this collection. It contained but fifty-six specimens of *Bibionidæ* and fifty *Dorilaidæ*. The majority of these have been adequately described and figured in other works (vide references at end of this paper) and are not being rediscussed. Keys to the genera and species are given in some of the cited references. All specimens were from the Belgian Congo and were collected by the Mission G. F. DE WITTE.

BIBIONIDÆ.

Seven species of *Bibionidæ* were present, these represent three genera.

Philia antipedalis ? (WIEDEMANN).

Dilophus antipedalis WIEDEMANN, 1818, Syst. Beschr. Eur. Zweifl., 1, p. 308.

Two female specimens are on hand which apparently belong to this species. The costa is, however, shorter than in other specimens of *antipedalis* which I have seen; the costa extends about one-fourth or one-fifth the distance between the apices of the third and fourth veins (the radial sector and vein M_{1+2}).

The specimens are from Lusinga, alt. 1.760 m, 11.VII to 8.XII.1947.

Philia erythræa (BEZZI).

Dilophus erythræus BEZZI, 1905, Firenze Boll. Soc. Ent. Ital., 37, pp. 205-206.

Seven specimens are in the collection from the following localities : Lusinga, alt. 1.760 m, 25.IV to 17.XII.1947 and Lusinga (riv. Dipidi), 12.VI.1945.

Philia nupta SPEISER.

Philia nupta SPEISER, 1914, Berl. Zeits. Deutsch. Ent. Ges., p. 1.

This was apparently the most common species of *Bibionidae* taken by this expedition. Thirty-six specimens are in the collection from the following localities : Mukana, alt. 1.810 m, 15.III.1948; riv. Kafwi, affl. dr. Lufwa, alt. 1.780 m, 5.III.1948; Lusinga, alt. 1.760 m, 12.III.1947 to 5.III.1948; Karibwe, affl. Lusinga, alt. 1.700 m, 8-10.III.1947 and Kamitungulu, affl. Lusinga, alt. 1.700 m, 4-7.III.1947.

Philia obsoleta HARDY.

Philia obsoleta HARDY, 1951, Journ. Kans. Ent. Soc., 24, pp. 91-92.

One female specimen from riv. Kilolomatembo, affl. Lusinga, alt. 1.750 m, 16.VII.1945.

Philia sp. ?
(Fig. 1.)

One female specimen is in the collection of a species, possibly undescribed, which is closely related to *P. nupta* SPEISER. The arrangement of the spines on the front tibiæ (fig. 1) is much like that of *nupta*, except that the pair of basad spines are located nearer the middle of the segment. The legs are predominantly black; only the front coxæ and trochanters and the basal portions of all the femora are rufous. The thorax is polished black, except for the rufous humeral ridges. The species is smaller than *nupta*.

Length : body, 4,2 mm; wings, 4,0 mm.

The specimen is from Lusinga, alt. 1.760 m, 1-8.XII.1947.

Bibio upembensis n. sp.
(Figs. 2a-b.)

This species is closely related to *Bibio afer* LOEW. It is distinguished by having vein M_{3+4} obsolete at the apex, not extending to the wing margin; by having nine segments in the antennæ (not ten); by the rufous or rufous tinged legs of the male and rufous coxæ, femora, sternum and pleura of the female; and by the yellow pile on the head and prothorax of the female.

Some specimens have a shortened $r-m$ crossvein and may run, imperfectly, to *B. hortulanus* (LINNAEUS), in couplet 2 of my key to the African *Bibio* (HARDY, 1950 b : 138). It is easily separated by the more fumose wings and by the incomplete vein M_{3+4} .

Male. — Head : Densely covered with brown to black pile. The antennæ are nine segmented, the last two are closely joined and appear as a single segment.




FIG. 1. — *Philia* n. sp. related to *nupta* SPEISER,
front tibia of female.

FIG. 2. — *Bibio upembensis* n. sp.
a. front tibia of male; b. hind tibia of female.

Thorax : Polished black except for the yellow humeral ridges. All of the pile is yellow. The halteres are dark brown to black.

Legs : The coxae, trochanters and tarsi are black; the femora and tibiæ are reddish brown, the hind femora are black at their bases. The femora and tibiæ of other specimens vary from all black, tinged with rufous, to all rufous. The inner spur on the front tibiæ is one-third to nearly one-half as long as the outer (fig. 2 a). The spurs of the hind tibiæ are moderately elongate and are blunt at apices (fig. 2 b).

Wings : Yellow to brownish fumose, darker along the costal margin. The costa ends at the apex of the radial sector. The anterior veins are brown. The posteriors are yellowish, slightly darker than the wing membrane. Vein M_{3+4} ends at a distance from the wing margin about equal to the length of the m crossvein. The $r-m$ crossvein is about one-third as long as the basal part of the radial sector. In some other specimens the $r-m$ is about half as long as the radial sector.

Abdomen : Subopaque black, densely covered with long yellow pile. The genitalia have been studied but do not appear to differ from those of other species of *Bibio*. No specific characters have been found in these structures.

Length : body, 7,0 mm; wings, 5,0 mm.

Female. — The thorax and abdomen, coxae and femora are all rufous except for the black central portion of the pronotum and the brown to blackish colored scutellum. All of the head and body pile is yellow, the tibiæ and tarsi are brown to black haired. The head is about one and one-fourth times longer than wide. The front is shining black and is slightly rugose. The front between the eyes is just slightly greater in width than the eyes. The eyes are about one and one-third times longer than that portion of the head behind the eyes.

Length : body, 7,5 mm; wings, 8,0 mm.

Holotype male : Dipidi, tête de source, alt. 1.700 m (rég. Lusinga), 17.VII.1947.

Allotype female : riv. Kilolomatembo, affl. Lusinga, alt. 1.750 m, 16.VII.1945. Seven paratypes, all males, from the following localities : same data as allotype; Lusinga, alt. 1.760 m, 3.VII.1947 and Parc National Albert : Ngesho, alt. 2.000 m, 3-6.IV.1934.

Type, allotype, and four paratypes returned to the « Institut des Parcs Nationaux du Congo Belge ». Two paratypes are in the U. S. National Museum and one is at the University of Hawaii.

Plecia zernyi HARDY.

Plecia zernyi HARDY, in press, Journ. Kans. Ent. Soc. (to appear in June, 1952).

One male specimen is in the collection from Lusinga, alt. 1.700 m, 17.III.1947.

DORILAIDÆ.

Eighteen species of *Dorilaïdæ* were present, representing three genera and two subgenera.

Dorilas (Eudorylas) abdominalis (LOEW).

Pipunculus abdominalis LOEW, 1857, Öfvers. K. Vet. Akad. Förhandl., 14, p. 374.

One specimen is in the collection from Mukana, near Lusinga, alt. 1.810 m, 28.V.1945.

Dorilas (Eudorylas) abruptus HARDY.

Dorilas (Eudorylas) abruptus HARDY, in press. *Bibionidæ et Dorilaïdæ* récoltés par F. J. FRANÇOIS dans le territoire de l'Urundi. Being published by the Institut royal des Sciences naturelles de Belgique.

One female specimen (a paratype) is in the collection from Lusinga, alt. 1.760 m, 1-8.XII.1947.

Dorilas (Dorilas) angustifacies HARDY.

Dorilas (Dorilas) angustifacies HARDY, 1949, Mém. Inst. roy. des Sc. nat. de Belg., 2^e sér., fasc. 36, p. 20.

One male is in the collection from Mubale, affl. g. de la Munte, alt. 1.480 m, 28.VI.1945.

Dorilas (Eudorylas) collarti HARDY.

Dorilas (Eudorylas) collarti HARDY, in press, Bull. Inst. roy. des Sc. nat. de Belg.

The female specimen in the collection has been designated as a paratype. It is from Lusinga, alt. 1.760 m, 1-8.VII.1947.

Dorilas (Dorilas) conspectus HARDY.

Dorilas (Dorilas) conspectus HARDY, 1949, Mém. Inst. roy. des Sc. nat. de Belg., 2^e sér., fasc. 36, pp. 27-28.

The propleural fan is rather poorly developed in this species and the hairs are often difficult to see, even when the propleura are plainly visible. The hairs are short and in some specimens only four to six hairs make up the fan.

Twelve specimens are in the collection from the following localities :

Lusinga, alt. 1.760 m, 8.IV.1947 to 2.V.1949; Mukana, près Lusinga, alt. 1.810 m, 1.VI.1945 and 14.IV.1947; Lusinga (riv. Dipidi), alt. 1.650 m, 12.VI.1945 and Lufwa, alt. 1.700 m, 16.III.1948.

Dorilas (Dorilas) damasi HARDY.

Dorilas (Dorilas) damasi HARDY, 1950, Expl. Parc Nat. Albert, Mission G. F. DE WITTE (1933-1935), fasc. 62, pp. 17-18.

Three specimens are in the collection from the following localities : riv. Kenia, affl. dr. Lusinga, alt. 1.585 m, 5-8.V.1949; Mukana, près de Lusinga, alt. 1.810 m, 20.VI.1945 and Lusinga, riv. Lusinga, alt. 1.810 m, 14.VI.1945.

Dorilas (Eudorylas) decorus HARDY.

Dorilas (Eudorylas) decorus HARDY, 1950, Expl. Parc Nat. Albert, Mission G. F. DE WITTE (1933-1935), fasc. 62, pp. 24-26.

Four specimens are in the collection from the following localities : Lusinga, alt. 1.760 m, 17.III to 6.XII.1947; Mukana, près de Lusinga, alt. 1.810 m, 28.V.1945 and Mubale, affl. g. de la Munte, 28.VI.1945.

Dorilas (Eudorylas) ghesquièrei HARDY.

Dorilas (Eudorylas) ghesquièrei HARDY, 1950, Expl. Parc Nat. Albert, Mission G. F. DE WITTE (1933-1935), fasc. 62, pp. 29-30.

One specimen is in the collection from Kaziba, alt. 1.140 m, 24.II.1948.

Dorilas (Eudorylas) mikenensis HARDY.

Dorilas (Eudorylas) mikenensis HARDY, 1950, Expl. Parc Nat. Albert, Mission G. F. DE WITTE (1933-1935), fasc. 62, pp. 33-34.

One female specimen is in the collection from Mukana, près de Lusinga, alt. 1.810 m, 28.V.1945.

Dorilas (Eudorylas) mutillatus (LOEW).


Pipunculus mutillatus LOEW, 1857, Öfver. K. Vet. Akad. Förhandl., 14, p. 374.

Three specimens are in the collection from the following localities : Karibwe, affl. Lusinga, alt. 1.700 m, 11.III.1947; Mubale, affl. g. de la Munte, alt. 1.480 m, 28.VI.1945 and Lusinga, alt. 1.760 m, 17.III.1947.

Dorilas (Dorilas) navus n. sp.

(Figs. 3a-d.)

This species is related to *D. visendus* HARDY and runs imperfectly to this species in couplet 21 of my key to the species known from the Belgian Congo (HARDY, 1950a : 9). The ninth segment is not visible from above and the membranous area is to the right side of the apex; not the left. The acutely pointed third antennal segment will also differentiate it, as will

FIG. 3. — *Dorilas (Dorilas) navus* n. sp.

- a. antenna;
- b. male genitalia, dorsal;
- c. male genitalia, right lateral;
- d. male genitalia, ventral.

the all yellow legs (except coxae) and the absence of strong bristles on the outside of the hind tibiae. In my key to the African *Dorilas* (HARDY, 1949a : 10) it runs imperfectly to *bellulus* HARDY. It differs by having the anterior crossvein at the basal third of the discal cell; the third costal section equal in length to the fourth and by the presence of a large membranous area covering all of the right, ventral portion of the apex of the male genitalia.

Male. — Head : The junction of the compound eyes is almost equal in length to the frontal triangle. The face is equal in width to the lower part of the front; the former is silvery pubescent, the latter is gray. The basal segments of the antennae are brown, the third is yellow and is acutely pointed below (fig. 3a). The occiput is entirely opaque gray, slightly brownish on the upper portion.

T h o r a x : Subshining black, tinged with yellow on the pleura. The dorsum is lightly brownish pollinose, the pleura are dusted with gray. The humeri and the knobs of the halteres are yellow, slightly tinged with brown. The propleura each have a fan of pale hairs.

L e g s : All yellow except for yellow-brown to black coxae. The femora are slightly swollen and flexor spines are moderately developed on the first two pairs. Apical spines are lacking on the tibiae and the hind pair have no strong erect bristles on the outside median surface.

W i n g s : Lightly brownish fumose with the stigma filling all of the third costal section. The third section is equal in length to the fourth and the two combined are slightly longer than the fifth. The anterior crossvein is situated at about the basal third of the discal cell and the last section of the fourth vein is slightly curved.

A b d o m e n : Chiefly polished black and slightly clavate in shape, broadest at the junction between the fourth and fifth segments. The first tergum has one or two bristles on each side, the remainder of the abdomen is nearly bare. The first tergum is entirely opaque brown to black. The second tergum is opaque brown except for the polished apical portion. The other terga are polished black with very narrow opaque bands across their basal margins.

G e n i t a l i a : About three-fourths as long as the fifth abdominal segment with the apical membranous area usually not visible in direct dorsal view (fig. 3 b). From a lateral view (fig. 3 c), or a ventral view, almost the entire right side is membranous. The ninth segment is brown, the claspers are yellow. The claspers are short and broad and each has a short finger-like lobe near the apex on the inner margin (fig. 3 d).

Length : body, 3,8 mm; wings, 4,4 mm.

Female unknown.

H o l o t y p e m a l e : Lusinga, alt. 1.760 m, 22-23.IV.1949. Two male paratypes, same data as type (one dated 16.III.1948).

The type and one paratype has been returned to the « Institut des Parcs Nationaux du Congo Belge ». One paratype has been deposited in the United States National Museum.

Dorilas (Dorilas) ruandensis HARDY.

Dorilas (Dorilas) ruandensis HARDY, 1950, Expl. Parc Nat. Albert, Mission G. F. DE WITTE (1933-1935), fasc. 62, pp. 21-22.

One specimen is present from Lusinga, alt. 1.760 m, 27.IV-2.V.1949.

Dorilas (Eudorylas) sinuosus HARDY.

(Fig. 4.)

Dorilas (Eudorylas) sinuosus HARDY, 1949, Mém. Inst. roy. des Sc. nat. de Belg., 2^e sér., fasc. 36, pp. 56-57.

Two male specimens of this unusual species are present in the collection, from Kalule Nord, rive g. face Mujinga-Kalenge, alt. 1.050 m, 28.II-3.III.1949 and Lusinga, alt. 1.760 m, 2-4.V.1949.

A few additional notes should be added to the original description of the male genitalia. The eighth segment is largely membranous. The membranous area extends over the median portion of the segment from near the anterior dorsal margin over the apex and around the venter to the anterior ventral margin (fig. 4). The eighth segment is longer than wide. The ninth segment is longer than wide and is scarcely longer than the claspers. The claspers are asymmetrical, the outer has a small finger-like lobe on its inner, subapical margin (fig. 4), the inner clasper is simple.

Dorilas (Dorilas) visendus HARDY.

Dorilas (Dorilas) visendus HARDY, 1950, Expl. Parc Nat. Albert, Mission G. F. DE WITTE (1933-1935), fasc. 62, pp. 23-24.

Two specimens are present from Lusinga, alt. 1.760 m, 2-4.V.1949.

Dorilas (Eudorylas) sp. ? poor cond.

One badly damaged specimen is present from Lusinga, alt. 1.760 m, 3.VII.1947.

Dorilas (subg. ?) n. sp. « A ».

(Fig. 5.)

One male specimen at hand appears to be undescribed but the prothorax is obscured by glue and the abdomen is slightly greased so it is not being named at this time.

The species runs to *D. congoensis* HARDY in my key to the *Dorilas* of the Belgian Congo (HARDY, 1950a : 10). It is distinguished by the narrow face and by the rufous coloration on the pleura and the first four abdominal segments; as well as by genital and wing characters. The abdomen is colored much as in *abdominalis* and related species but the abdomen is opaque to faintly shining and not polished. The narrow face and the rufous coloration on the abdomen would suggest that the species might be related to *D. conspectus* HARDY but the abdomen is more extensively rufous and is not polished; the legs are all yellow; the anterior crossvein is situated at the basal fourth of the discal cell and the last section of the fourth vein is

strongly curved. The dorsal aspects of the genitalia are as in figure 5. A small basal cleft is present on the extreme right side. This does not show from a direct dorsal view.

One male is present from Lusinga (riv.), alt. 1.810 m, 9.VI.1945.




FIG. 4. — *Dorilas (Eudorylas) sinuosus* HARDY, male genitalia, ventral view.

FIG. 5. — *Dorilas* n. sp. « A », male génitalia, dorsal.

FIG. 6. — *Wittella candidula* (HARDY), male genitalia, ventral.

FIG. 7. — *Wittella lusingensis* n. sp.

a. antenna; b. male genitalia, dorsal; c. male genitalia, ventral
(from a paratype specimen).

Wittella lusingensis n. sp.

(Figs. 7a-c.)

This species is very closely related to *W. candidula* (HARDY). It is best distinguished by the dark brown to black antennæ; the rather thickly pilose thorax and abdomen and by the differences in the male genitalia as shown in figures 6 and 7c.

Male. — **Head** : The compound eyes are joined on the front for a distance about equal to nearly one-half of the length of the frontal triangle. The frontal triangle and the face are gray pubescent, subshining black in ground color. The upper portion of the front, below the vertex, is polished black. The upper third of the occiput is polished black, the remainder is gray pollinose. The antennæ are dark brown to black. The third segment is small and is obtuse, rounded below (fig. 7a).

Thorax : Entirely polished black except for the yellow humeri and halteres. Rather conspicuously yellow pilose on the margins, scutellum and down the dorsocentral areas. The propleura have a fan of hairs.

Legs : Entirely yellow except for the black coxae and brown apical subsegments of the tarsi. The flexor spines are moderately developed on the middle femora. The hind tibiæ have three or four strong erect bristles near the middle on the outside surface. The front and middle tibiæ have rather conspicuous apical spines. The femora are all slender.

Wings : Faintly brownish fumose. The third costal section is about one-third as long as the fourth and the fourth section is about equal in length to the fifth. The stigma fills all of the third costal section. The anterior crossvein is situated at the basal fifth of the discal cell. The last section of the fourth vein is slightly curved.

Abdomen : Entirely polished black, rather thickly covered with erect yellow hairs. The abdomen is slightly clavate, it is broadest at the junction of segments four and five. The first tergum has a row of long yellow bristles on each side.

Genitalia : From a dorsal view the genitalia are about one-half as long as the fifth abdominal segment and a large membranous area is present at the apex of the eighth segment (fig. 7b). From a ventral view the ninth segment is longer than wide and two or more times longer than the claspers. The claspers are asymmetrical in shape, the inner is much broader than the outer. The inner clasper is approximately as broad as long (fig. 7c). (Note : the information on the ventral aspects of the genitalia is taken from a paratype specimen.)

Length : body, 4,0 mm; wings, 4,5 mm.

Female unknown.


Holotype male : Lusinga, alt. 1.760 m, 12-17.XII.1947. Two paratypes, males : Kalumengongo, alt. 1.780 m, 18.IV.1947 and Mukana, près de Lusinga, alt. 1.810 m, I-IV.1947.

The type and one paratype have been returned to the « Institut des Parcs Nationaux du Congo Belge ». One paratype has been deposited in the United States National Museum.

Tömösváryella mbuyensis n. sp.

(Figs. 8a-e.)

This species appears more closely related to *T. apicalis* HARDY than to any other known species. It is distinguished by the keel-like development of the underside of the hind trochanters; by the short junction of the compound eyes of the male on the front, and by the differences in the male genitalia as shown in the figures.

FIG. 8. — *Tömösváryella mbuyensis* n. sp.

- a. antenna;
- b. hind trochanter of male;
- c. male genitalia, dorsal;
- d. male genitalia, ventral;
- e. female abdomen, lateral.

In my key to the African *Tömösváryella* (HARDY, 1949a : 64-66), this species runs to *apicalis* HARDY in couplet 12. In my key to the *Tömösváryella* of the Belgian Congo (HARDY, 1950a : 42-43) it runs to *congoana* HARDY, in couplet 6. It is distinguished from both of these species by the characters given above.

Male. — **Head** : The eyes are joined for just a short distance on the front. The junction is about equal in length to six or seven of the eye facets. The frontal triangle is gray pubescent, the portion of the front

below the vertex is shining black. The face is equal in width to the broadest portion of the front. The antennæ are dark brown to black, the third segment is short acuminate (fig. 8 a). The labellum is yellow.

T h o r a x : Subshining black in ground color, brownish pollinose on the dorsum, gray on the sides. The humeri and halteres are yellow.

L e g s : Predominantly black, the apices of the tibiæ and femora, the bases of the tibiæ and the basal subsegments of the tarsi are yellow. The hind trochanters have a ridge-like development on the underside, this is raised into a slight point (fig. 8 b). The hind trochanters have rather long setæ below.

W i n g s : Hyaline, faintly iridescent. The third costal section is about one-half as long as the fourth and the two sections combined are less than half as long as the fifth. The anterior crossvein is situated near the middle of the discal cell and the last section of the fourth vein is straight or nearly so. The last section of the fifth vein is shorter than the posterior crossvein.

A b d o m e n : Subshining black in ground color, dusted with grayish brown pollen on the dorsum, gray on the sides. The sides are nearly straight, it is slightly broadest at the apex of the third segment. The abdomen is rather sparsely covered with suberect hairs.

G e n i t a l i a : About three-fourths as long as the fifth abdominal segment and with a moderately large membranous area at its apex (fig. 8 c). This membranous area is over half as broad as the eighth segment. The ninth segment is slightly longer than wide and about equal in length to the claspers. The claspers are slender and rather elongate, the inner is slightly longer than the outer; they are both blunt pointed (fig. 8 d).

Length : body and wings, 2,8-3,0 mm.

F e m a l e . — The front is expanded in the middle and at this point is wider than the face. The lower half of the face is grey, the upper half is shining black. The hind trochanters are not ridged as in the male and are covered with gray pubescence on the underside. The ovipositor is slender and straight, the piercer is one and one-half to two times longer than its base and extends to about the base of segment two (fig. 8 e). The piercer is rufous in color, the base is subshining black, the latter is oblong in shape.

Length : body, 2,4 mm; wings, 2,6 mm.

H o l o t y p e m a l e : Buye-Bala, affl. g. Muye, alt. 1.750 m, 25-31.III.1948. Allotype, same data as type. Paratypes, three males : Two same locality as type; 1 specimen, same data; 1 specimen, 1-7.IV.1948 and one from Mukana, près Lusinga, alt. 1.810 m, 15-19.I.1948.

The type, allotype and one paratype have been returned to the « Institut

des Parcs Nationaux du Congo Belge ». One paratype has been deposited in the U. S. National Museum collection, and one is at the University of Hawaii.

Tömösváryella subvirescens (LOEW).

Pipunculus subvirescens LOEW, 1872, Berl. Ent. Zeitschr., 16, p. 87.

Four specimens are in the collection from the following localities : Lusinga, alt. 1.760 m, 22-23.IV.1949 and Kalule Nord, rive g. face Mujinga-Kalenge, alt. 1.050 m, 28.II-3.III.1949.

REFERENCES CITED.

- HARDY, D. E., 1949a. The African *Dorilaidæ* (*Mém. Inst. roy. des Sci. nat. de Belgique*, 2^e série, fasc. 36, 1-80, 6 pls.).
- 1949b. New *Dorilaidæ* from the Belgian Congo [*Bull. Inst. roy. des Sci. nat. de Belgique*, 25 (39), 1-10, 15 figs.).
- 1950a. Exploration du Parc National Albert, Miss. G. F. de Witte (1933-1935), fasc. 62 : *Dorilaidæ*, 1-53, 7 pls.
- 1950b. A monographic study of the African *Bibionidæ*. Part I : Introduction and genus *Bibio* (*Journ. Kans. Ent. Soc.*, 23, 137-153, 13 figs.).

University of Hawaii,
Department of Zoology and Entomology.

TABLE DES MATIÈRES

	Pages
PLECOPTERA :	
1. Perlidæ	5
COLEOPTERA :	
2. Histeridæ	13
3. Chrysomelidæ	23
4. Scolytoidea	49
DIPTERA :	
5. Bibionidæ and Dorilaidæ	57

AVIS

L'Institut des Parcs Nationaux du Congo Belge a commencé, en 1937, la publication des résultats scientifiques des missions envoyées aux Parcs Nationaux, en vue d'en faire l'exploration.

Les divers travaux paraissent sous forme de fascicules distincts. Ceux-ci comprennent, suivant l'importance du sujet, un ou plusieurs travaux d'une même mission. Chaque mission a sa numérotation propre.

Les fascicules peuvent s'acquérir séparément.

L'Institut des Parcs Nationaux du Congo Belge n'accepte aucun échange.

BERICHT

Het Instituut der Nationale Parken van Belgisch Congo heeft in 1937 de publicatie aangevangen van de wetenschappelijke uitslagen der zendingen welke naar de Nationale Parken afgevaardigd werden, ten einde ze te onderzoeken.

De verschillende werken verschijnen in vorm van afzonderlijke afleveringen welke, volgens de belangrijkheid van het onderwerp, één of meer werken van dezelfde zending bevatten. Iedere zending heeft haar eigen nummering.

De afleveringen kunnen afzonderlijk aangeschaft worden.

Het Instituut der Nationale Parken van Belgisch Congo neemt geen ruilingen aan.

FASCICULES PARUS

HORS SÉRIE :

Les Parcs Nationaux et la Protection de la Nature.

Discours prononcé par le Roi Albert à l'installation de la Commission du Parc National Albert.

Discours prononcé par le Duc de Brabant à l'African Society, à Londres, à l'occasion de la Conférence Internationale pour la Protection de la Faune et de la Flore africaines.

La Protection de la Nature. Sa nécessité et ses avantages, par V. VAN STRAELEN, 1937.

VERSCHEENEN AFLEVERINGEN

BUITEN REEKS :

De Nationale Parken en de Natuurbescherming.

Redevoering uitgesproken door Koning Albert op de vergadering tot aanstelling der Commissie van het Nationaal Albert Park.

Redevoering door den Hertog van Brabant gehouden in de African Society, te Londen, bij de gelegenheid van de Internationale Conferentie voor de Bescherming van de Afrikaansche Fauna en Flora.

De Natuurbescherming. Haar noodzakelijkheid en haar voordeelen, door V. VAN STRAELEN, 1937.

Exploration du Parc National Albert. — Exploratie van het Nationaal Albert Park

I. — Mission G. F. DE WITTE (1933-1935).

I — Zending G. F. DE WITTE (1933-1935).

Fasc.	Afl.		
1.	G. F. DE WITTE (Bruxelles), <i>Introduction</i>		1937
2.	C. ATTEMS (Vienne), <i>Myriopodes</i>		1937
3.	W. MICHAELSEN (Hamburg), <i>Oligochäten</i>		1937
4.	J. H. SCHUURMANS STEKHOVEN Jr (Utrecht), <i>Parasitic Nematoda</i>		1937
5.	L. BURGEON (Tervueren), <i>Carabidae</i>		
	M. BANNINGER (Giessen), <i>Carabidae (Scaritini)</i>		1937
6.	L. BURGEON (Tervueren), <i>Lucanidae</i>		1937
7.	L. BURGEON (Tervueren), <i>Scarabaeidae (S. Fam. Cetoniinae)</i>		1937
8.	R. KLEINE (Stettin), <i>Brenthidae und Lycidae</i>		1937
9.	H. SCHOUTEDEN (Tervueren), <i>Oiseaux</i>		1938
10.	S. FRECHKOP (Bruxelles), <i>Mammifères</i>		1938
11.	J. BEQUAERT (Cambridge, Mass.), <i>Vespides solitaires et sociaux</i>		1938
12.	A. JANSSENS (Bruxelles), <i>Onitini (Coleoptera Lamellicornia, Fam. Scarabaeidae)</i>		1938
13.	L. GSCHWENDTNER (Linz), <i>Haliplidae und Dytiscidae</i>		1938
14.	E. MEYRICK (Marlborough), <i>Pterophoridae (Tortricina and Tineina)</i>		1938
15.	C. MOREIRA (Rio de Janeiro), <i>Passalidae</i>		1938
16.	R. J. H. TEUNISSEN (Utrecht), <i>Tardigraden</i>		1938
17.	W. D. HINCKS (Leeds), <i>Dermoptera</i>		1938
18.	R. HANITSCH (Oxford), <i>Blattids</i>		1938
19.	G. OCHS (Frankfurt a. Main), <i>Gyrinidae</i>		1938
20.	H. DEBAUCHE (Louvain), <i>Geometridae</i>		1938
21.	A. JANSSENS (Bruxelles), <i>Scarabaeini (Coleoptera Lamellicornia, Fam. Scarabaeidae)</i>		1938
22.	J. H. SCHUURMANS STEKHOVEN Jr et R. J. H. TEUNISSEN (Utrecht), <i>Nématodes libres terrestres</i>		1938
23.	L. BURGEON (Tervueren), <i>Curculionidae, S. Fam. Apioninae</i>		1938
24.	M. POLL (Tervueren), <i>Poissons</i>		1939
25.	A. JANSSENS (Bruxelles), <i>Oniticellini (Coleoptera Lamellicornia, Fam. Scarabaeidae)</i>		1939
26.	L. BURGEON (Tervueren), <i>Histeridae</i>		1939
27.	<i>Arthropoda : Hexapoda : 1. Orthoptera : Mantidae, par M. BEIFER (Wien); 2. Gryllidae, par L. CHOPARD (Paris); 3. Coleoptera : Cicindelidae, par W. HORN (Berlin); 4. Rutelinae, par F. OHAUS (Mainz); 5. Heteroceridae, par R. MAMITZA (Wien); 6. Prioninae, par A. LAMEERE (Bruxelles); Arachnoidea : 7. Opiliones, par C. FR. ROEWER (Bremen)</i>		1939
28.	A. HUSTACHE (Lagny), <i>Curculionidae</i>		1939
29.	A. JANSSENS (Bruxelles), <i>Coprini (Coleoptera Lamellicornia, Fam. Scarabaeidae)</i>		1940
30.	L. BERGER (Bruxelles), <i>Lepidoptera-Rhopalocera</i>		1940
31.	V. LABOTSSIÈRE (Paris), <i>Galerucinae (Coleoptera Phytophaga, Fam. Chrysomelidae)</i>		1940
32.	V. LALLEMAND (Bruxelles), <i>Homoptera (Cicadidae, Cercopidae, Fulgoridae, Dictyophoridae, Ricaniidae, Cixiidae, Derbidae, Flatidae)</i>		1941
33.	G. F. DE WITTE (Bruxelles), <i>Batractiens et Reptiles, avec Introduction de V. VAN STRAELEN</i> .		1941

Fasc.	
Afl.	
34. L. MADER (Wien), <i>Coccinellidae</i> . — I. Teil	1941
II. Teil	1950
35. R. PAULIAN (Paris), <i>Aphoditinae</i> (<i>Coleoptera Lamellicornia</i> , Fam. <i>Scarabaeidae</i>)	1942
36. A. VILLIERS (Paris), <i>Languriinae et Cladoxeninae</i> (<i>Coleoptera Clavicornia</i> , Fam. <i>Erotylidae</i>)	1942
37. L. BURGEON (Tervueren), <i>Chrysomelidae</i> (S. Fam. <i>Eumolpinae</i>)	1942
38. A. JANSSENS (Bruxelles), <i>Dynastinae</i> (<i>Coleoptera Lamellicornia</i> , Fam. <i>Scarabaeidae</i>) ...	1942
39. V. LABOISSIÈRE (Paris), <i>Halticinae</i> (<i>Coleoptera Phytophaga</i> , Fam. <i>Chrysomelidae</i>) ...	1942
40. F. BORCHMANN (Hamburg), <i>Lagriidae und Alleculidae</i>	1942
41. H. DEBAUCHE (Louvain), <i>Lepidoptera Heterocera</i>	1942
42. E. UHMANN (Stollberg), <i>Hispaniae</i>	1942
43. <i>Arthropoda : Arachnoidea</i> : 1. <i>Pentastomida</i> , par R. HEYMONS (Berlin); <i>Hexapoda</i> : 2. <i>Orthoptera</i> : <i>Phasmidae</i> , par K. GUENTHER (Dresden); 3. <i>Hemiptera : Membracidae</i> , by W. D. FUNKHOUSE (Lexington U.S.A.); 4. <i>Coleoptera : Silphidae</i> , par A. JANSSENS (Bruxelles); 5. <i>Dryopidae</i> , par J. DELÈVE (Bruxelles); 6. <i>Lymexylonidae</i> , par L. BURGEON (Tervueren); 7. <i>Bostrichidae</i> , par P. LESNE (Paris); 8. <i>Scarabaeidae : Geotrupinae</i> , par A. JANSSENS (Bruxelles); 9. <i>Cassidinae</i> , von A. SPAETH (Wien); 10. <i>Ipidae</i> , von H. EGGER (Bad Nauheim); 11. <i>Platypodidae</i> , par K. E. SCHEDL (Hann. Münden); 12. <i>Hymenoptera : Sphegidae</i> , by G. ARNOLD (Bulawayo) ...	1943
44. G. MARLIER (Bruxelles), <i>Trichoptera</i>	1943
45. H. SCHOUTEDEN (Tervueren), <i>Reduviidae, Emesidae, Henicocephalidae (Hemiptera Heteroptera)</i>	1944
46. R. PAULIAN (Paris), <i>Hybosoridae et Trogidae</i> (<i>Coleoptera Lamellicornia</i>) ...	1944
47. H. DE SAEGER (Bruxelles), <i>Microgasterinae (Hymenoptera Apocrita)</i> ...	1944
48. G. SCHMITZ (Louvain), <i>Chalcididae (Hymenoptera Chalcidoidea)</i> ...	1946
49. H. DEBAUCHE (Louvain), <i>Mymaridae (Hymenoptera Apocrita)</i> ...	1949
50. H. DE SAEGER (Bruxelles), <i>Euphoriniae (Hymenoptera Apocrita, Fam. Braconidae)</i> ...	1946
51. A. COLLART (Bruxelles), <i>Helomyzinae (Diptera Brachycera, Fam. Helomyzidae)</i> ...	1946
52. P. VANSCHUYTBROECK (Bruxelles), <i>Sphaerocerinae (Diptera Acalyptratae, Fam. Sphaeroceridae)</i> ...	1948
53. H. DE SAEGER (Bruxelles), <i>Cardiochilinae, Sigalphinae (Hymenoptera Apocrita, Fam. Braconidae)</i> ...	1948
54. A. THÉRY (Neuilly), <i>Buprestidae (Coleoptera Sternoxia)</i> ...	1948
55. M. GOETGHEBUER (Gand), <i>Ceratopogonidae (Diptera Nematocera)</i> ...	1948
56. H. SCHOUTEDEN (Tervueren), <i>Coreidae (Hemiptera Heteroptera)</i> ...	1948
57. H. F. STROHECKER (Miami), <i>Endomychidae (Coleoptera Clavicornia)</i> ...	1949
58. R. POISSON (Rennes), <i>Hémiptères aquatiques</i> ...	1949
59. M. CAMERON (London), <i>Staphylinidae (Coleoptera Polyphaga)</i> ...	1950
60. J. PASTEEELS (Bruxelles), <i>Tenthredinidae (Hymenoptera Tenthredinoidea)</i> ...	1949
61. F. C. FRASER (Bournemouth), <i>Odonata</i> ...	1949
62. D. ELMO HARDY (Honolulu, Hawaii), <i>Dorilaidæ (Diptera)</i> ...	1950
63. J. BALFOUR-BROWNE (London), <i>Palpicornia</i> ...	1950
64. R. LAURENT, <i>Genres Afrixalus et Hyperolius (Amphibia Salientia)</i> ...	1950
65. D. ELMO HARDY (Honolulu, Hawaii), <i>Bibionidae (Diptera Nematocera)</i> ...	1950
66. J. VERBEKE (Gand), <i>Sciomyzidae (Diptera Cyclorrhapha)</i> ...	1950
67. H. OLDROYD (London), <i>Genera Hæmatopota and Hippocentrum (Diptera, Fam. Tabanidae)</i> ...	1950
68. A. REICHENSPERGER (Bonn), <i>Paussidæ</i> ...	1950
69. H. HAUPt (Halle), <i>Pompilidæ (Hymenoptera Sphecoidea)</i> ...	1950
70. <i>Hexapoda</i> : 1. <i>Orthoptera : Tridactylidæ</i> , par L. CHOPARD (Paris); 2. <i>Hemiptera : Coccidæ</i> , par P. VAYSSIÈRE (Paris); 3. <i>Coleoptera : Trogositidæ</i> , par G. FAGEL (Bruxelles); <i>Erotylidæ</i> , von K. DELGESKAMP (Berlin); <i>Bostrichidæ</i> , par J. VRYDAGH (Bruxelles); <i>Megalopodinæ</i> , by G. E. BRYANT (London); <i>Anthribidæ</i> , by K. JORDAN (Tring); 4. <i>Diptera : Therevidæ</i> , par P. VANSCHUYTBROECK (Bruxelles); <i>Conopidæ</i> , par P. VANSCHUYTBROECK (Bruxelles); 5. <i>Hymenoptera : Chrysididæ</i> , von S. ZIMMERMANN (Wien) ...	1950
71. K. ERMISCH (Radiumbad), <i>Mordellidæ (Coleoptera Heteromera)</i> ...	1950
72. J. VERBEKE (Gand), <i>Taxiapterinæ (Diptera Cyclorrhapha, Fam. Micropezidæ)</i> ...	1951
73. P. L. G. BENOIT (Tervueren), <i>Dryinidæ (Hymenoptera Aculeata); Evanidæ (Hymenoptera Terebrantia)</i> ...	1951
74. P. VANSCHUYTBROECK (Bruxelles), <i>Dolichopodidae (Diptera Brachycera Orthorrhapha)</i> .	1951
75. N. BRUCE (Stockholm), <i>Cryptophagidae (Coleoptera Polyphaga)</i> ...	1951
76. M. C. MEYER (Orono), <i>Hirudinea</i> ...	1951
77. 1. <i>Thysanoptera</i> , by H. PRIESTER (Cairo); 2. <i>Suctoria (Aphaniptera)</i> , par J. COOREMAN (Bruxelles); 3. <i>Homoptera</i> , par V. LALLEMAND et H. SYNAVE (Bruxelles); 4. <i>Coleoptera : Sagridæ</i> , par P. JOLIVET (Bruxelles); <i>Clytridæ</i> , par P. JOLIVET (Bruxelles); 5. <i>Diptera : Asilidæ</i> , by S. W. BROMLEY (Stamford, U.S.A.); <i>Sinuliidæ</i> , g. <i>Simulium</i> , by P. FREEMAN (London) ...	1951
78. J. VERBEKE (Zürich), <i>Psilidæ (Diptera Cyclorrhapha)</i> ...	1952

II. — Mission H. DAMAS (1935-1936).

II. — Zending H. DAMAS (1935-1936).

Fasc.

Afl.

- | | | |
|-----|-----------------------------------------------------------------------------------------------|----------------------------|
| 1. | H. DAMAS (Liège), <i>Recherches Hydrobiologiques dans les Lacs Kivu, Édouard et Ndalaga</i> | 1937 |
| 2. | W. ARNDT (Berlin), <i>Spongilliden</i> | 1938 |
| 3. | P. A. CHAPPUIS (Cluj, Roumanie), <i>Copépodes Harpacticoides</i> | 1938 |
| 4. | E. LELOUP (Bruxelles), <i>Moerisia Alberti nov. sp. (Hydropolyte dulcicole)</i> | 1938 |
| 5. | P. DE BEAUCHAMP (Strasbourg), <i>Rotifères</i> | 1939 |
| 6. | M. POLL (Tervueren), avec la collaboration de H. DAMAS (Liège), <i>Poissons</i> | 1939 |
| 7. | V. BREHM (Eger), <i>Cladocera</i> | 1939 |
| 8. | F. HUSTEDT (Ploen), <i>Süßwasser Diatomeen</i> | 1949 |
| 9. | J. H. SCHUURMANS STEKHOVEN Jr (Utrecht), <i>Nématodes libres d'eau douce</i> | 1944 |
| 10. | J. H. SCHUURMANS STEKHOVEN Jr (Utrecht), <i>Nématodes parasites</i> | 1944 |
| 11. | G. MARLIER (Bruxelles), <i>Trichoptera</i> | 1943 |
| 12. | W. KLIE (Bad Pyrmont), <i>Ostracoda</i> | 1944 |
| 13. | G. MARLIER (Bruxelles), <i>Collemboles</i> | 1944 |
| 14. | J. COOREMAN (Bruxelles), <i>Acaris</i> | 1948 |
| 15. | A. ARCANGELI (Torino), <i>Isopodi terrestri</i> | 1950 |
| 16. | F. GUIGNOT (Avignon), <i>Dytiscidae et Gyrinidae (Coleoptera Adephaga)</i> | 1948 |
| 17. | H. BERTRAND (Dinard), <i>Larves d'Hydrocanthares</i> | 1948 |
| 18. | O. LUNDBLAD (Stockholm), <i>Hydrachnellae</i> | 1949 |
| 19. | W. CONRAD (Bruxelles), P. FRÉMY (St.-Lô) et A. PASCHER (Prague), <i>Algues et Flagellates</i> | 1949 |
| 20. | M.-L. VERRIER (Paris), <i>Éphéméroptères</i> | 1951 |
| 21. | FR. KIEFER (Konstanz-Staad), <i>Copépodes</i> | (Sous presse.) (Ter pers.) |

III. — Mission P. SCHUMACHER (1933-1936).

III. — Zending P. SCHUMACHER (1933-1936).

1. P. SCHUMACHER (Antwerpen), *Die Kivu-Pygmaen und ihre soziale Umwelt im Albert-Nationalpark* 1943
 2. P. SCHUMACHER (Antwerpen), *Anthropometrische Aufnahmen bei den Kivu-Pygmaen*. 1939

IV. — Mission J. LEBRUN (1937-1938).

IV. — Zending J. LEBRUN (1937-1938).

V. — Mission S. FRECHKOP (1937-1938).

V. — Zending S. FRECHKOP (1937-1938).

- | | |
|---------------------------------------------------------------------------------------------------|------|
| 1. S. FRECHKOP (Bruxelles), <i>Mammifères</i> | 1943 |
| 2. R. VERHEYEN (Bruxelles), <i>Oiseaux</i> | 1947 |

VI. — Missions J. VERHOOGEN (1938 et 1940).

VI. — Zendingen J. VERHOOGEN (1938 en 1940).

1. J. VERHOOGEN (Bruxelles), *Les éruptions 1938-1940 du volcan Nyamuragira* 1948

Vol

FLORE DES SPERMATOPHYTE DU PARC NATIONAL ALBERT

Exploration du Parc National Albert et du Parc National de la Kagera. Exploratie van het Nationaal Albert Park en van het Nationaal Park der Kagera.

- I. — Mission L. VAN DEN BERGHE (1936). I. -- Zending L. VAN DEN BERGHE (1936).

Fasc.
Afl.

1. L. VAN DEN BERGHE (Anvers), *Enquête parasitologique*. — I. — *Parasites du sang des vertébrés*
 2. L. VAN DEN BERGHE (Anvers), *Enquête parasitologique*. — II. — *Helminthes parasites*.

1942
1943

Exploration du Parc National de la Kagera. — Exploratie van het Nationaal Park der Kagera.

- I. — Mission J. LEBRUN (1937-1938). I. — Zending J. LEBRUN (1937-1938).

1. J. LEBRUN, L. TOUSSAINT, A. TATON (Bruxelles), *Contribution à l'étude de la flore du Parc National de la Kagera ...*

1948

- ## II. — Mission S. FRECHKOP (1938).

- ## II. — Zending S. FRECHKOP (1938).

1. S. FRECHKOP (Bruxelles), *Mammifères*
 2. R. VERHEYEN (Bruxelles), *Oiseaux*

1944

Exploration du Parc National de l'Upemba. — Exploratie van het Nationaal Upemba Park.

- I. — Mission G. F. DE WITTE en collaboration avec
W. ADAM, A. JANSENS, L. VAN MEEUWEN et R. VERHEYEN
(1946-1949).

- I. — Zending G. F. DE WITTE in medewerking met
W. ADAM, A. JANSSENS, L. VAN MEEL en R. VERHEYEN
(1946-1949).

Fasc.
Afl.

Exploration des Parcs Nationaux du Congo Belge — Exploratie der Nationale Parken van Belgisch Congo.

- I. — Mission H. HEDIGER - J. VERSCHUREN (1948).

- I. — Zending H. HEDIGER - J. VERSCHUREN (1948).

1. II. HEDIGER (Bâle), *Observations sur la psychologie animale dans les Parcs Nationaux du Congo Belge*

1951

ASPECTS DE VEGETATION DES PARCS NATIONAUX DU CONGO BELGE

**VEGETATIEBEELDEN
DER NATIONALE PARKEN VAN BELGISCH CONGO**

AVIS

Les Aspects de Végétation des Parcs Nationaux du Congo Belge paraissent par fascicules de six planches, accompagnées de notices explicatives.

La publication est divisée en séries, consacrées chacune à un *Parc National du Congo Belge*.

La première série a pour objet le *Parc National Albert*.
Les fascicules peuvent s'acquérir séparément.

L'Institut des Parcs Nationaux du Congo Belge n'accepte aucun échange

FASCICULES PARUS

SÉRIE 1. — PARC NATIONAL ALBERT.

Volume I.

PUBLICATIONS SÉPARÉES

LOSSE PUBLICATIES

Mammifères et Oiseaux protégés au Congo Belge, par S. FRECHKOP, avec Introduction de V. VAN STRAelen (Épuisé.) (Uitgeput.)
Contribution à l'étude de la Morphologie du Volcan Nyamuragira, par R. HOIER (Rutshuru) 1939
Intimaux protégés au Congo Belge et dans le Territoire sous mandat du Ruanda-Urundi, ainsi que les espèces dont la protection est assurée en Afrique (y compris Madagascar) par la Convention Internationale de Londres du 8 novembre 1933 pour la protection de la Faune et de la Flora africaines, avec la Législation concernant la Chasse, la Pêche, la Protection de la Nature et les Parcs Nationaux au Congo Belge et dans le Territoire sous Mandat du Ruanda-Urundi, par S. FRECHKOP, en collaboration avec G. F. DE WITTE, J.-P. HARROY et E. HUBERT, avec Introduction de V. VAN STRAelen (1941).
 (Épuisé.) (Uitgeput.)

Beschermd Dieren in Belgisch Congo en in het Gebied onder mandaat van Ruanda-Urundi, evenals de Soorten waarvan de bescherming verzekerd is in Afrika (met inbegrip van Madagascar) door de Internationale Overeenkomst van Londen van 8 November 1933 voor de bescherming van de Afrikaansche Flora en Fauna, met de Wetgeving betreffende de Jacht, de Visscherij, de Natuurbescherming en de Nationale Parken van Belgisch Congo en in het Gebied onder mandaat van Ruanda-Urundi, door S. FRECHKOP, in medewerking met G. F. DE WITTE, J.-P. HARROY en E. HUBERT, met Inleiding van V. VAN STRAELEN (1943) (Épuisé.) (Uitgeput.)

Les territoires bioécoégraphiques du Parc National Albert par W. BORYNS (Épuisé.) (Uitgeput.) 1948

Les territoires biogéographiques du Parc National Albert, par W. ROBYNS 1948

A travers plaines et volcans au Parc National Albert, par R. HOIER 1950

Parcs Nationaux du Congo Belge ... 1949

Contribution à l'étude éthologique des mammifères du Parc National de l'Upemba, par R. VERHEYEN. 1951

animaux protégés au Congo Belge et dans le Territoire sous mandat du Rwanda-Urundi, édition 1952

BERICHT

De Vegetatiebeelden der Nationale Parken van Belgisch Congo verschijnen in afleveringen van zes platen, van verklarende aantekeningen vergezeld.

De publicatie is ingedeeld in reeksen, waarvan elke aan één der Nationale Parken van Belgisch Congo gewijd is.

De eerste reeks handelt over het *Nationaal Albert Park*. De afleveringen kunnen afzonderlijk aangeschaft worden.

Het Instituut der Nationale Parken van Belgisch Congo neemt geen ruilingen aan.

VERSCHENEN AFLEVERINGEN

REEKS I. — NATIONAAL ALBERT PARK.

Boekdeel I.

Afl. 1-2 — W. ROBYNS (Brussel), *Algemeen overzicht der vegetatie* (volgens de fotografische documentatie der zending G. F. DE WITTE) ... 1937